



India's **Most Comprehensive** & the **Most Relevant**
Test Series designed according to the latest pattern of exams!



JEE MAIN



JEE ADV.



BITSAT



WBJEE



MHT CET

and many more...

[Click here to join Test Series for 2022](#)

It's time for you to crack upcoming IIT JEE Main & Advanced and other competitive exams with India's Most Trusted Online Test Series. **Many questions at JEE Main 2021 were same/similar to the ones asked in our test series.** That's the power of our test series!

Trusted by **thousands of students**
& **their parents** across the nation

Our result in JEE Main 2021

150+

Got **99+ percentile** (overall)

301

Got **99+ percentile** in one
or more subjects

85%

Improved their score by
25 percentile

89%

Felt **overall confident** after
the test series

[Click here to join Test Series for 2022](#)



FREE Question Bank & Previous Year Questions for



JEE MAIN



JEE ADV.



BITSAT



WBJEE



MHT CET

and many more...



Click on this button



🤖 Why download MARKS?

- 📖 Chapter-wise PYQ of JEE Main, JEE Advanced, NEET, AIIMS, BITSAT, WBJEE, MHT CET etc.
- 📖 Chapter-wise NTA Abhyas questions
- 🎯 Daily practice challenge and goal completion
- 📌 Bookmark important questions and add them to your notebooks
- 👉 Create unlimited Custom Tests

And all this for **FREE**. Yes, **FREE**! So what are you waiting for, download MARKS now.

4.8

Rating on Google Play

30,000+

Students using daily

1,00,000+

Questions available



Click on this button

Differentiation

Derivative

- (i) The rate of change of one quantity with respect to some other quantity has a great importance. For example the rate of change of displacement of a particle with respect to time is called its velocity and rate of change of velocity is called its acceleration. The rate of change of a quantity 'y' with respect to another quantity 'x' is called the derivative or differential coefficient of y with respect to x .
- (ii) Let $y = f(x)$ be a continuous function of an independent variable x. Let δx be an arbitrary small change in the value of x and δy be the corresponding change in that of y. Then limit of the ratio $\frac{\delta y}{\delta x}$ as $\delta x \rightarrow 0$, if exists, is named as the derivative or differential coefficient of x with respect to y and it is denoted by $\frac{dy}{dx}$

$$\text{So } \frac{dy}{dx} = \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x}$$

$$\text{i.e. } \frac{dy}{dx} = \lim_{\delta x \rightarrow 0} \frac{f(x + \delta x) - f(x)}{\delta x} = f'(x)$$

Derivative of y with respect to x is also denoted by the following symbols : $y \phi$, y_1 , Dy

The process of finding derivative of a function is called differentiation.

Derivatives of some standard functions

Some Differentiation Formulae

$$(i) \quad \frac{d}{dx} (\text{constant}) = 0$$

$$(iii) \quad \frac{d}{dx} (e^x) = e^x$$

$$(v) \quad \frac{d}{dx} (\log_e x) = \frac{1}{x}$$

$$(vii) \quad \frac{d}{dx} (\sin x) = \cos x$$

$$(ix) \quad \frac{d}{dx} (\tan x) = \sec^2 x$$

$$(xi) \quad \frac{d}{dx} (\sec x) = \sec x \tan x$$

$$(xiii) \quad \frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}} ; -1 < x < 1$$

$$(xv) \quad \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2} ; x \in \mathbb{R}$$

$$(xvii) \quad \frac{d}{dx} (\sec^{-1} x) = \frac{1}{|x|\sqrt{x^2-1}} , |x| > 1$$

$$(ii) \quad \frac{d}{dx} (x^n) = nx^{n-1}$$

$$(iv) \quad \frac{d}{dx} (a^x) = a^x \log_e a$$

$$(vi) \quad \frac{d}{dx} (\log_a x) = \frac{1}{x \log_e a}$$

$$(viii) \quad \frac{d}{dx} (\cos x) = -\sin x$$

$$(x) \quad \frac{d}{dx} (\cot x) = -\operatorname{cosec}^2 x$$

$$(xii) \quad \frac{d}{dx} (\operatorname{cosec} x) = -\operatorname{cosec} x \cot x$$

$$(xiv) \quad \frac{d}{dx} (\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}} ; -1 < x < 1$$

$$(xvi) \quad \frac{d}{dx} (\cot^{-1} x) = -\frac{1}{1+x^2} , \forall x \in \mathbb{R}$$

$$(xviii) \quad \frac{d}{dx} (\operatorname{cosec}^{-1} x) = \frac{-1}{|x|\sqrt{x^2-1}} ; |x| > 1$$

$$(xix) \frac{d}{dx} (e^{ax} \sin bx) = e^{ax} (a \sin bx + b \cos bx) = \sqrt{a^2 + b^2} e^{ax} \sin (bx + \tan^{-1} \frac{b}{a})$$

$$(xx) \frac{d}{dx} (e^{ax} \cos bx) = e^{ax} (a \cos bx - b \sin bx) = \sqrt{a^2 + b^2} e^{ax} \cos (bx + \tan^{-1} \frac{b}{a})$$

$$(xxi) \frac{d}{dx} |x| = \frac{x}{|x|} \quad (x \neq 0)$$

$$(xxii) \frac{d}{dx} \log |x| = \frac{1}{x} \quad (x \neq 0)$$

Some theorems of differentiation

$$(i) \frac{d}{dx} (f(x) \pm g(x)) = \frac{d}{dx} (f(x)) \pm \frac{d}{dx} (g(x))$$

$$(ii) \frac{d}{dx} (k f(x)) = k \frac{d}{dx} (f(x)), \text{ where } k \text{ is any constant}$$

$$(iii) \frac{d}{dx} (f_1(x) \cdot f_2(x)) = (f_1(x)) \frac{d}{dx} (f_2(x)) + (f_2(x)) \frac{d}{dx} (f_1(x))$$

$$(iv) \frac{d}{dx} \left(\frac{f_1(x)}{f_2(x)} \right) = \frac{f_2(x) \frac{d}{dx} (f_1(x)) - f_1(x) \frac{d}{dx} (f_2(x))}{(f_2(x))^2}$$

(v) Derivative of the function of the function .

If 'y' is a function of 't' and 't' is a function of 'x', then

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}$$

$$\text{Thus } \frac{d}{dx} \{g[f(x)]\} = \frac{dg\{f(x)\}}{df(x)} \cdot \frac{d}{dx} f(x) = g' \{f(x)\} \cdot f'(x)$$

(vi) Derivative of parametric equation

If $x = \phi(t)$, $y = \psi(t)$ then

$$\frac{dy}{dx} = \frac{dy}{dt} \bigg/ \frac{dx}{dt}$$

(vii) Derivative of a function w.r.t another function

If $f(x)$ and $g(x)$ are two function of a variable x , then

$$\frac{d(f(x))}{d(g(x))} = \frac{d(f(x))}{dx} \bigg/ \frac{d(g(x))}{dx}$$

Differentiation of Implicit Functions

If in an implicit function $f(x, y) = 0$, y cannot be expressed in terms of x , then we differentiate both sides of the given equation w.r.t. x and collect all terms containing $\frac{dy}{dx}$ on L.H.S.

NOTE : In the above process we obtain dy/dx in terms of both x and y . If we want dy/dx in terms of x only, then let us first express y in terms of x .

Short method of differentiation for implicit functions

If $f(x, y) = \text{constant}$, then

$$\frac{dy}{dx} = -\left(\frac{\partial f}{\partial x}\right) / \left(\frac{\partial f}{\partial y}\right)$$

where $\left(\frac{\partial f}{\partial x}\right)$ and $\left(\frac{\partial f}{\partial y}\right)$ are partial derivatives of $f(x, y)$ with respect to x and y respectively.

[By partial derivative of $f(x, y)$ with respect to x , we mean the derivative of $f(x, y)$ with respect to x when y is treated as a constant.]

Logarithmic differentiation

If differentiation of an expression or an equation is done after taking log on both sides, then it is called logarithmic differentiation. Generally we apply this method when given expression is in one of the following forms

- (i) $[f(x)]^{g(x)}$
- (ii) product of three or more function

Some Suitable substitutions

- (i) If the function involve the term $\sqrt{a^2 - x^2}$, then substitute $x = a \sin \theta$ or $x = a \cos \theta$
- (ii) If the function involve the term $\sqrt{x^2 + a^2}$, then substitute $x = a \tan \theta$ or $x = a \cot \theta$
- (iii) If the function involve the term $\sqrt{x^2 - a^2}$, then substitute $x = a \sec \theta$ or $x = a \operatorname{cosec} \theta$
- (iv) If the function involve the term $\left(\sqrt{\frac{a-x}{a+x}}\right)$, then substitute $x = a \cos \theta$ or $x = a \cos 2\theta$

Differentiation of a Determinant Function

$$\text{If } F(x) = \begin{vmatrix} f & g & h \\ l & m & n \\ u & v & w \end{vmatrix}$$

Where $f, g, h, l, m, n, u, v, w$ are functions of x and differentiable, then

$$F'(x) = \begin{vmatrix} f' & g' & h' \\ l & m & n \\ u & v & w \end{vmatrix} + \begin{vmatrix} f & g & h \\ l' & m' & n' \\ u & v & w \end{vmatrix} + \begin{vmatrix} f & g & h \\ l & m & n \\ u' & v' & w' \end{vmatrix}$$

$$\text{or } F'(x) = \begin{vmatrix} f' & g & h \\ l' & m & n \\ u' & v & w \end{vmatrix} + \begin{vmatrix} f & g' & h \\ l & m' & n \\ u & v' & w \end{vmatrix} + \begin{vmatrix} f & g & h' \\ l & m & n' \\ u & v & w' \end{vmatrix}$$