



India's **Most Comprehensive** & the **Most Relevant**  
Test Series designed according to the latest pattern of exams!



JEE MAIN



JEE ADV.



BITSAT



WBJEE



MHT CET

and many more...

[Click here to join Test Series for 2022](#)

It's time for you to crack upcoming IIT JEE Main & Advanced and other competitive exams with India's Most Trusted Online Test Series. **Many questions at JEE Main 2021 were same/similar to the ones asked in our test series.** That's the power of our test series!

Trusted by **thousands of students**  
& **their parents** across the nation

Our result in JEE Main 2021

**150+**

Got **99+ percentile** (overall)

**301**

Got **99+ percentile** in one  
or more subjects

**85%**

Improved their score by  
**25 percentile**

**89%**

Felt **overall confident** after  
the test series

[Click here to join Test Series for 2022](#)



# FREE Question Bank & Previous Year Questions for



JEE MAIN



JEE ADV.



BITSAT



WBJEE



MHT CET

and many more...



Click on this button



## 🤖 Why download MARKS?

- 📖 Chapter-wise PYQ of JEE Main, JEE Advanced, NEET, AIIMS, BITSAT, WBJEE, MHT CET etc.
- 📖 Chapter-wise NTA Abhyas questions
- 🎯 Daily practice challenge and goal completion
- 📌 Bookmark important questions and add them to your notebooks
- 👉 Create unlimited Custom Tests

And all this for **FREE**. Yes, **FREE**! So what are you waiting for, download MARKS now.

4.8

Rating on Google Play

30,000+

Students using daily

1,00,000+

Questions available



Click on this button

# Probability

---

## 1. EXPERIMENT:

An operation which results in some well-defined outcomes is called an experiment.

## 2. RANDOM EXPERIMENT:

The experiment about which no confirm prediction is possible, is called the random experiment.

e.g. (i) Tossing a coin, (ii) Throwing a die etc.

## 3. SAMPLE SPACE:

The set of all possible outcomes of an experiment is called the sample space, which is denoted by  $S$ .

e.g. (i) On throwing a die, sample space is

$$S = \{1, 2, 3, 4, 5, 6\} \text{ and } n(S) = 6$$

(ii) On tossing two coins, sample space is

$$S = \{H, T\} \times \{H, T\} = \{HH, HT, TH, TT\}.$$

$$n(S) = 4$$

## 4. EVENT:

Every subset of sample space is defined as the event. It is denoted by  $E$ .

e.g. (i) The event of appearing odd numbers in throwing a die is

$$E = \{1, 3, 5\}$$

$$\text{and } E \subseteq S, \text{ where } S = \{1, 2, 3, 4, 5, 6\}$$

## 5. TYPES OF EVENTS :

(i) Null Event or Impossible Event: It has no member and is denoted by  $\phi$ .

e.g. (a) On throwing a die the event of appearing member 7 is an impossible event.

(ii) Sure Event: Since  $S \subseteq S$ , so  $S$  is an event which is called sure event.

(iii) Simple Event or Elementary Event: It has only one member.

e.g. (b) On throwing a die the event of appearing prime even number is  $E = \{2\}$  which is a simple event.

(iv) Mixed Event or Composite event or

Compound Event: This event has two or more than 2 members.

e.g.: On throwing a die the event of appearing even numbers is

$$E = \{2, 4, 6\} \text{ which is a compound event.}$$

(v) Mutually Exclusive Events: Events  $E_1$  and  $E_2$  are mutually exclusive if  $E_1 \cap E_2 = \phi$

e.g.: In drawing a card from a pack of 52 cards, the events

$A \rightarrow$  the event when the card drawn is spade

$B \rightarrow$  the event when the card drawn is a heart are mutually exclusive events.

(vi) Mutually Exclusive and Exhaustive Events:

Events  $E_1, E_2, E_3, \dots, E_n$  are mutually exclusive and exhaustive events if

(a)  $E_i \cap E_j = \phi$ , where  $i \neq j$  and  $i, j = 1, 2, 3, \dots, n$ .

(b)  $E_1 \cup E_2 \cup E_3 \cup \dots \cup E_n = S$

e.g. On throwing a die if

$A$  = the event of appearing odd numbers =  $\{1, 3, 5\}$

$B$  = The event of appearing even numbers =  $\{2, 4, 6\}$

Then  $A \cap B = \phi$

and  $A \cup B = S = \{1, 2, 3, 4, 5, 6\}$

$\therefore A$  and  $B$  are mutually exclusive and exhaustive events.

(v) Complement of an Event  $E$ :

It is denoted by  $E'$  or  $E^C$  or  $\bar{E}$  and it is the set of all sample points of sample space not of event  $E$ .

On throwing a die,  $S = \{1, 2, 3, 4, 5, 6\}$

If event be  $E = \{1, 2, 3\}$

then  $\bar{E} = \{4, 5, 6\}$

Note : (a)  $A \cup \bar{A} = S$  (b)  $A \cap \bar{A} = \phi$ .

## 6. PROBABILITY OF AN EVENT:

$$P(E) = \frac{n(E)}{n(S)} = \frac{\text{no. of favourable cases}}{\text{no. of favourable and infavourable cases}}$$

Remarks:

- $P(S) = 1$
- $P(\phi) = 0$
- $0 \leq P(E) \leq 1$
- If  $A \subseteq B$  then  $P(A) \leq P(B)$
- $P(A) + P(\bar{A}) = 1$

## 7. ODDS IN FAVOUR OF EVENT E:

$$\begin{aligned} \text{Odds in favour of event } E &= \frac{P(E)}{P(E')} = \frac{P(E)}{1 - P(E)} \\ &= a : b, \text{ say.} \end{aligned}$$

$$\text{If odds in favour of event } E \text{ be } a : b, \text{ then } P(E) = \frac{a}{a + b}$$


## 8. ODDS AGAINST EVENT E:

$$\text{Odds against event } E = \frac{P(E')}{P(E)} = \frac{1 - P(E)}{P(E)} = c : d, \text{ say}$$

It odds against event E be  $c : d$ , then

$$P(E) = \frac{d}{c+d}.$$

## 9. IDEAS ABOUT A PACK OF 52 CARDS:



## 10. SOME OTHER TYPES OF EVENTS:

**Independent Events:** Two or more events are said to be independent if occurrence or non-occurrence of any one of them does not affect the occurrence or non-occurrence of other.

e.g. If a bag contains 3 red and 2 black balls and if two balls are drawn one by one with replacements, then events are independent events.

**Dependent Events:** Two or more events are said to be dependent, if occurrence or non-occurrence of any one of them affects occurrence or non-occurrence of others.

e.g. Let a bag contains 3 red and 2 black balls. Two balls are drawn one by one without replacement. Then these events are dependent events.

**Equally Likely Events:** Outcomes are said to be equally likely which we have no reason to believe that one is more likely to occur than the other.

e.g. When an unbiased die is thrown all the six faces 1, 2, 3, 4, 5, 6 are equally likely to come up.

## 11. ADDITION THEOREM FOR PROBABILITY:

(i) If A and B the any two events, then  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

(ii) If A and B are mutually exclusive events, then  $P(A \cup B) = P(A) + P(B)$

(iii) If A, B, C are any three vents, then

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$$

(iv) If A, B and C are mutually exclusive events, then

$$P(A \cup B \cup C) = P(A) + P(B) + P(C)$$

$$(v) \quad P(A \cup B) \leq P(A) + P(B)$$

Some other results:

$$(i) \quad P(\bar{A} \cap B) = P(B) - P(A \cap B) = P(B - A)$$

$$(ii) \quad P(A \cap \bar{B}) = P(A) - P(A \cap B) = P(A - B)$$

$$(iii) \quad P\{(A \cap \bar{B}) \cup (\bar{A} \cap B)\} = P(A) + P(B) - 2P(A \cap B)$$

$$(iv) \quad P(\bar{A} \cap \bar{B}) = 1 - P(A \cup B)$$

$$(v) \quad P(\bar{A} \cup \bar{B}) = 1 - P(A \cap B)$$

(vi) If A, B, C be three events the

(a) P(atleast two of A, B, C occur)

$$= P(A \cap B) + P(B \cap C) + P(C \cap A) - 2P(A \cap B \cap C)$$

(b) P(exactly two of A, B, C occur) =  $P(A \cap B) + P(B \cap C) + P(A \cap C) - 3P(A \cap B \cap C)$

(c) P(exactly one of A, B, C, occurs)

$$= P(A) + P(B) + P(C) - 2P(A \cap B) - 2P(B \cap C) - 2P(A \cap C) - 3P(A \cap B \cap C)$$

## 12. CONDITIONAL PROBABILITIES

Let A and B be any two events such that  $B \neq \phi$  or  $n(B) \neq 0$  or  $P(B) \neq 0$ , then  $P(A/B)$  denotes the conditional probability of occurrence of event A when B has been already occurred.

e.g. A coin is tossed thrice. If E be the event of showing atleast two heads and F the event of showing head in the first throw, then

$$P(E/F) = \frac{P(E \cap F)}{P(F)} = \frac{3/8}{4/8} = \frac{3}{4}$$

## 13. MULTIPLICATION THEOREM:

(i) If A and B be any two events then

$$P(A \cap B) = P(A) \cdot P(B/A)$$

$$= P(B) \cdot P(A/B)$$

(ii) If events A and B are independent events, then

$$P(A \cap B) = P(A) \cdot P(B) = P(AB)$$

## 14. BAYE'S THEOREM:

Let S be the sample space and let  $E_1, E_2, \dots, E_n$  be n mutually exclusive and exhaustive events associated with a random experiment. If A is any event which occurs with  $E_1$  or  $E_2$  or .....  $E_n$ , then

$$P(E_i/A) = \frac{P(E_i) \times P(A/E_i)}{P(E_1).P(A/E_1) + P(E_2).P(A/E_2) + \dots + P(E_n).P(A/E_n)} \text{ for } n = 1, 2, 3, \dots, n.$$

**15. BINOMIAL PROBABILITY DISTRIBUTION:**

If for a sample space  $S$ ,  $n$  = total no. of trial

$p$  = probability of event of success for one - trial

$q$  = probability of event of failure for one trial

so that  $p + q = 1$ , then probability for  $r$  success trial is

$$P(X = r) = {}^nC_r \cdot p^r \cdot q^{n-r}.$$

**16. MEAN AND VARIANCE OF BINOMIAL DISTRIBUTION:**

Mean =  $np$  and Variance =  $npq$

and mean > variance

Also standard deviation =  $\sqrt{npq}$

