

India's Most Comprehensive & the Most Relevant Test Series designed according to the latest pattern of exams!











JEE MAIN

JEE ADV.

WBJEE

MHT CET

and many more...

Click here to join Test Series for 2022

It's time for you to crack upcoming IIT JEE Main & Advanced and other competitive exams with India's Most Trusted Online Test Series. Many questions at JEE Main 2021 were same/similar to the ones asked in our test series. That's the power of our test series!

Trusted by thousands of students & their parents across the nation

Our result in JEE Main 2021

150+

Got 99+ percentile (overall)

301

Got **99+ percentile** in one or more subjects

85%

Improved their score by **25 percentile**

89%

Felt **overall confident** after the test series

Click here to join Test Series for 2022



FREE Question Bank & Previous Year Questions for



JEE MAIN










JEE ADV.

BITSAT W

WBJEE MHT CET

and many more...





Why download MARKS?

- Chapter-wise PYQ of JEE Main, JEE Advanced, NEET, AIIMS, BITSAT, WBJEE, MHT CET etc.
- Chapter-wise NTA Abhyas questions
- Taily practice challenge and goal completion
- Bookmark important questions and add them to your notebooks
- Create unlimited Custom Tests

And all this for FREE. Yes, FREE! So what are you waiting for, download MARKS now.

4.8

Rating on Google Play

30,000+

Students using daily


1,00,000+

Questions available




KINEMATICS-1D

- Kinematics in which we deal with the motion i.e. change in position with respect to a either fixed system or a
 moving system (frame of reference) without taking into account of its cause or force.
- For analysing motion of a particle, there are some physical quantities as its parameters
 - (i) Position or position vector which is the location of the particle with respect to system of co-ordinate axes.
 - (ii) Displacement the change in position vector.
 - (iii) Velocity The time rate of change of displacement
 - (iv) Acceleration The time rate of change of velocity.



In figure (i) motion is along positive x-direction $\stackrel{\text{u.u.}}{AB} = \text{displacement}$ in time $\Delta t = \Delta x$




In fig (ii) displacement in time Δt , $AB = OB - OA = \Delta r$ = change in position vector.

In any kind of motion, the magnitude of displacement is less than or equal to distance (actual path traversed).

• $AB = \Delta S = Distance$ covered in time Δt . AB = Displacement is time $\Delta t = \Delta r$ or path length

$$\frac{\Delta S}{\Delta t} = \text{Average speed in time } \Delta t = \frac{\text{Distance covered}}{\text{Time elapsed}}$$

$$\frac{\Delta r}{\Delta t}$$
 = Average velocity in time $\Delta t = \frac{Displacement}{Time elapsed}$



If for different interval the particle has same average velocity then motion is said to be in uniform motion or otherwise motion is said to be in non-uniform motion and if particle is in non-uniform motion then velocity at a

[2] Kinematics

point is defined as $\lim_{\Delta t \to 0} \frac{\Delta r^r}{\Delta t} = \frac{dr^r}{dt} = \frac{r}{v}$ called instantaneous velocity, for one dimension $\frac{r}{v} = \frac{dr^r}{dt}$. The instantaneous velocity at any print is equal to average velocity in any interval for uniform motion i.e., $\frac{r}{v} = \frac{dr^r}{dt}$.

As
$$\Delta S \ge |\Delta_r^1|$$

$$\frac{\Delta S}{\Delta t} \ge \left| \frac{\Delta r}{\Delta t} \right| \text{ i.e. Av. speed } \ge \mid \text{Av. velocity}|.$$

Equality sign holds when motion is in one-dimensional and unidirectional.

But for $\Delta t \to 0$, $\frac{\Delta s}{\Delta t} = \frac{|\Delta_r^r|}{\Delta t} \Rightarrow \frac{ds}{dt} = \frac{|dr|}{dt}$ i.e. magnitude of instantaneous velocity is the speed at a point.

• If force is in the direction of motion then motion is accelerated and if force is opposite to direction of motion then it is deaccelerated or retared. Rate of change of velocity w.r. to time is same for any interval and it is equal to instantaneous acceleration. Motion is said to be uniformly accelerated otherwise non-uniformly accelerated.

$$\frac{r}{a_{\rm av}} = \frac{\Delta \overline{v}}{\Delta t} = \mbox{Average acceleration in time interval } \Delta t \; . \label{eq:average}$$

$$\lim_{\Delta t \to 0} \frac{\Delta v^{\Gamma}}{\Delta t} = \frac{dv^{\Gamma}}{dt} = \text{acceleration at a point} = \text{constant for uniformally accelerated}$$

$$\frac{d^{\Gamma}}{dt}$$
 = f(t) or f(s) of f(v) then motion is said to be non-uniformly accelerated.

• Translational kinematical equation of motion for uniformly accelerated motion.

$$\overset{\mathbf{r}}{\mathbf{S}_{n}} = \overset{\mathbf{r}}{\mathbf{u}} + \frac{1}{2}\overset{\mathbf{r}}{\mathbf{a}}(2n-1)$$
 (S_n = Displacement is in last one second.)

• Equation for non uniformly accelerated motion.

$$\frac{d\vec{x}}{dt} = \vec{r}$$


$$\frac{d\vec{v}}{dt} = f(t) \text{ or } f(s) \text{ or } f(v) \text{ and } \frac{d\vec{v}}{ds} \cdot \vec{r} = f(s)$$

- Direction of velocity is in direction of change of displacement while direction of acceleration is in direction of change in velocity.
- Vertical motion under gravity Motion of a particle under the effect of gravity This is motion in one dimension with constant acceleration $a = g = 9.8 \text{ ms}^{-2} = 32 \text{ ft sec}^2$.

If a particle is projected veritcally up with velocity u then time of ascent = $\frac{u}{g}$ and during which height attained

$$H = \frac{u^2}{2g} \text{ . Time of descent} = \frac{u}{g} \qquad \qquad |\overset{\mathbf{r}}{v_A}|(\uparrow) = |\overset{\mathbf{r}}{v_B}|(\downarrow)$$

Kinematics [3]



 $t_{AL} = t_{LB}$ where L is the point up to which particle ascends. where A and B are at same height from surface of earth.


• Effect of air resistance on motion of a particle in vertical motion under gravity - Air resistance is a force that acts on the body opposite to direction of motion and its value depends on the speed.


KINEMATICAL GRAPH -


Displacement time graph

From displacement time graph we find


- (i) slope at a point gives the instantaneous velocity
- (ii) slope of the chord gives the average velocity in the given interval x-t curve for constant acceleration.







VELOCITY TIME CURVE -





From velocity time graph we have


- (i) slope of the tangent at a point gives the acceleration at a point.
- (ii) slope of the chord gives the average acceleration in a given interval.
- (iii) Area under the v-t curve gives displacement, taking area positive for upper of time axis and area negative for lower of time axis.
- (iv) Area under the v-t curve gives distance when all the areas either above or below t-axis taken to be positive.

[4] Kinematics

Velocity time curve for uniformly accelerated motions.







Acceleration time curve -

It is of very physical significance in which area under the curve in the given interval is the change in velocity in that interval.